

PEG LINKERS

PEGylation is the process of attaching PEG linkers to molecules such as a drug, protein or vesicle. Due to its high water-solubility and non-immunogenicity, PEG has been widely used in a pharmaceutical & biotech R&D to improve drug solubility and stability, optimize DMPK and discover new delivery formats.

BroadPharm's PEG Linker represents a specific, single pure molecule with high purity. These monodisperse PEGs are designed with defined PEG lengths and various functional groups.

Over 2,000 PEG reagents are supplied by BroadPharm.



PEG Linkers (in stock)			
Alkyne PEG	Bromo PEG	Non-PEG linker	PEG Tosylate
Amino PEG	Cleavable PEG	PEG Acid	PEG-X-PEG
Aminoxy PEG	DBCO PEG	PEG Aldehyde	Poly PEG
BCN-PEG	DNP-PEG	PEG azide	Propargyl PEG
Benzyl-PEG	DSPE PEG	PEG Maleimide	Sugar PEG
Biotin PEG	Fluorescent PEG	PEG NHS ester	TCO-PEG
Bis-PEG-acid	Fmoc PEG	PEG Peptide	Tetrazine-PEG
Bis-PEG-NHS	Hydroxy PEG	PEG PFP ester	Thiol PEG
Boc-PEG	m-PEG	PEG Phosphonate	
Branched PEG	MeNH-PEG	PEG Sulfonic acid	



重松貿易株式会社 化学品部

Shigematsu & Co., Ltd.

<大阪本社>

〒541-0047 大阪市中央区淡路町 2 丁目 2 番 5 号

TEL : (06)6231-6146 FAX : (06)6231-6149

<つくばライフサイエンスオフィス>

〒300-2655 茨城県つくば市万博公園西 F26 街区 1 インプレス 103

TEL : (029)848-3220 FAX : (029)848-3221

WEB : www.shigematsu-bio.com/ Mail : info@shigematsu-bio.com



